

DESCRIPTION OF HEALTH RISKS ASSOCIATED WITH THE USE OF ILLICIT DRUGS AND ABUSE OF ALCOHOL

Controlled Substances - Uses & Effects							
Classification	Drug Name	Common or Brand	Dependence Physical	Potential Psycho- logical	Acute Effects	Health Risks and Effects of Long-term Use*	Overdose Effects
ALCOHOL	Ethanol	Beer Wine Distilled Liquor	High High High	High High High	Lowered Inhibitions Impaired Judgement, Vision Loss of Motor Skills, Coordination Sturred Speech	Hypertension Liver Damage Cardiovascular Disease Toxic Psychosis Neurologic Damage Addiction with Severe Withdrawal	Coma Possible Death
NARCOTICS	'Opium Morphine Codeine 'Heroin Meperidine Other	¹ Paregoric, Dover S Power ¹ Morphine ¹ Tylenol W/Codeine, Robitussin Heroin ¹ Demerol, Pethadol ² Dilaudid, Lentine, Percodan	High High High High High High	High High High High High High	Euphoria, Drowsiness Respiratory Depression Constricted Pupils Nausea, Vomit Analgesia (Pain Relief)	Loss of Appetite Constipation Risk of AIDS & Hepatitis from I.V. Drug Use	Shallow Perspiration Clammy Skin Convulsions, Corna Possible Death May be Toxic if Mixed with Alcohol
DEPRESSANTS	Chloral Hydrate Barbiturates Garnma Hydroxybutric Acid Garmma Butyrolactone Glutethimide Ketarnine Methaqualone Tranquilizers	*Noctec, Somnos *Nembutal, Phonobarbitol, Seconal, Tuinol Date Rape Drug, Liquid Eestacy, Cherrymeth, GBL *Doriden Special K, K *Optimil, Parest, Quaalude, Sopor *Equanil, Valium, Tranxene	Moderate High Moderate-High Unknown High Unknown High Moderate	Moderate High High Unknown High Unknown High High	Surred Speech Disorientation Drunken Behavior Without the Odor of Alcohol Euphoria Lowared Inhibitions Loss of Motor Skills Blackouts Pelavation, Depression Hallucinations	Addiction with Withdrawal Toxic Psychosis Weak, Rapid Pulse Hallucinations Naussa Seizures Insomnia Anxiety Tremors Dizziness Loss of Peripheral Vision	Shallow Respiration Cold, Clammy Skin Blackout Coma Possible Death May be Toxic if mixed with Alcohol
STIMULANTS	'Cocaine 'Amphetamine Phenmetrazine Methylphenidate Other Methylenedioxy- methamphetamine	Serax, Xanax, Rohyphal, Dalmane, Dormate, Placidyl, Valmid Cocaine Hydrochloride *Benzedrine, Biphetamine, Desoxyn, Dexedrine *Preludin *Ritalin *Cylert, Didrex, Pre-Sate, Sanorax, Voranil Ecstacy, MDMA	Possible Possible Possible Possible Possible	High High High High High	Feeling of Well-Being Excitation, Euphoria Increased Alertness Increased Blood Pressure, Pulse Insomnia, Appetite Loss	Delusions, Hallucinations (Toxic Psychosis) Possible Organ/Tissue Damage	High Agitation Temperature Increase Hallucinations Convulsions Heart Attack, Stroke High Blood Pressure Loss of Consciousness Saizures
HALLUNCINO- GENICS	*LSD *Mescaline *Psilocybin *MDA *PCP	Acid, Lysergic Acid Peyote, Peyote Buttons, Mescalitor Magic Mushrooms, Shrooms Angel Dust, Crystal, Cernylan (Veterinary)	Low Low Low Unknown Unknown	Unknown Unknown Unknown Unknown Unknown	Dilated Pupils Increased Blood Pressure, Pulse Deliusions/Hallucinations Distorted Perception of Time, Space, and Reality	Hallucinogens, Especially LSD, may Intensify Existing Psychosis Possible Violent Behavior	Intense Bad Trips Psychosis Possible Sudden Death
INHALANTS	Airplane Glue Lighter Fluid Aerosols Spray Paints Paint Thinner Gasoline Poppers	Active Ingredient: Toluene Active Ingredient: Naphalene Active Ingred: Chloroplourocarbon Active Ingred: Petroleum Distillate Active Ingred: Petroleum Distillate Active Ingred: Petroleum Distillate Active Ingred: Petroleum Jistillate Active Ingred: Petroleum Jistillate	Unknown Unknown Unknown Unknown Unknown Unknown Unknown Unknown	Unknown Unknown Unknown Unknown Unknown Unknown Unknown	Slurred Speech Drunken Behavior Impaired Judgement Poor Coordination	Hallucinations Possible Darnage to Bone Marrow, Lungs, Kidneys, Liver, Heart, Brain, Eyes	Unconsciousness Coma Possible Toxic Reaction Possible Sudden Death
CANNABIS	'Marijuana 'Hashish 'Hash Oil	Grass, Pot, Weed, Dope Hash	Low Low Low	Moderate Moderate Moderate	Tachycardia, Reddened Eyes Euphoria, Profound Humor Altered Time/Space Percepton Short-Torm Memory Loss Increased Appetite	Cardiovascular Damage as with Smoking Tobacco	Insomnia, Hyperactivity Panic Attacks, Paranoia Possible Toxic Reaction if Treated w/other Chemicals
TOBACCO	Nicotine	Cigarettes Cigars	Moderate Moderate	High High	Relaxation Stimulation	Cardiovascular Disease Respiratory Illness	Possible Death at very High Dosage Levels

^{*} Alcohol and other drug use during pregnancy increases risk of physical harm to fetus

² Psychoactive drug effects refer to use at a greater than prescribed therapeutic dosage level

Additional risk of harm from toxic impurities in street drugs